



Post translational modification of proteins in the context of immune recognition



F. Hudecz^{1,2}

¹ Research Group of Peptide Chemistry, Hungarian Academy of Sciences,
Eötvös Loránd University (ELTE),

² Department of Organic Chemistry, Institute of Chemistry, ELTE, Budapest, Hungary


In vivo post-translational modifications - diversity

alkyl	Cleavage of peptide bond
acyl (O-,N-,S-)	N-terminals Met or fMet
N-terminal,	signalpeptide
Lys,	precursor activation (proinsulin → insulin)
Ser, Thr	
amide	Disulphid bond formation
C-terminal	
phosphoric acid ester (Ser, Thr, Tyr)	Isomerisation (Pro, Asp)
sulphonic acid ester	
glycosylation	Coupling of nucleotide (e.g. flavine)
O- in Golgi (Ser, Thr)	
N- in RER (Asn)	
nitrosation	
desamidation	
decarboxylation	
Arg desamination,	
citrullination (Arg → citrullin)	
hydroxylation (Pro, Lys)	
oxidation	
gamma-carboxylation (e.g. Glu)	
beta-elimination (e.g. Thr → alkene)	
	Coupling of protein/peptide :
	sumoylation (SUMO protein)
	ubiquitination (ubiquitin)
	neddylation (Nedd)

SUMOylation (Small Ubiquitin-like Modifier)

Involved in nuclear-cytosolic transport,
transcriptional regulation,
apoptosis
protein stability,
but, not in degradation

- SUMO proteins: 100 aa. , 12 kDa, 4 isoforms
- Post-translational modification
- Activation: - cleavage of 4 residues at the C-terminal
- Attachment to target protein by using three enzymes .



R. Geiss-Friedlander & F. Melchior
Nature Rev. Mol. Cell Biol. 8, 947-956 (2007)

Ubiquitination




KUNGL.
VETENSKAPSAKADEMIEN
THE ROYAL SWEDISH ACADEMY OF SCIENCES



Nobel Prize in Chemistry, 6 October 2004
A. Ciechanover, A. Hershko, I. Rose



- Ubiquitin: protein (76 aa., 8.5 kDa)
- Almost all tissues of eukaryotic organisms
- „Label“ the degradation of the attached protein by transporting to proteasome.
- Isopeptide linkage (4)
- Enzymes involved:
 - E1 (ub activation)
 - E2 (ub conjugation to ϵ -amino group (Lys), thiol (Cys) by thioester, and to OH (Thr/Ser) by ester formation)
 - E3 (ub ligation)



NEDDylation

(Neural-precursor-cell-expressed developmentally down-regulated)

- Function:
activation/regulation of ubiquitin




Post-translational modification: Immune recognition related diseases

Modification	Autoantigen	Disease
Acetylation	Myelin basic protein	Multiple sclerosis
Citrullination	Collagen type II Myelin basic protein	RA Multiple sclerosis
Deamidation	Insulin	Type I diabetes
Glycosylation	Insulin proceptor Collagen type II Thyrotropin receptor Myelin oligodendrocyte glycoprotein Mucin glycoprotein (MUC2)	Diabetes RA Graves disease MS Colon carcinoma
Isoaspartylation	snRNP	Systemic lupus erythematosus
Lipoylation	PDC-E2	Primary biliary cirrhosis
Phosphorylation	Myelin basic protein	Multiple sclerosis
Methylation	Sm, D1,D3	Systemic lupus erythematosus
Transglutamination	Histone H2	Systemic lupus erythematosus
Tyrosine nitration	Mitochondrial proteins	Experimental autoimmune uveitis


Posttranslational modification of Auto-antigens p.41
in "Autoantibodies" Eds. Y. Shoenfeld, M. E. Gershwin, P.- L. Meroni, pp. 838, (2007) Elsevier

Post-translational modification: influence on immune recognition


glycosylation




citrullination




deamidation




disulfide




phosphorylation




sulfurylation




nitration




hydroxylation




isomerisation



methylation/alkylation




acetylation/acylation





The effect of post-translational modification on T-cell immune recognition: Isomerisation of Asp to β -Asp

T cell response



structure of horse heart cytochrome c (PDB:1HRC)




Assay:

- B10.A mice immunized with 100 μg peptide 94-104 with CFA
- after 10 days cell suspension from lymph nodes
- antigen stimulation with peptide a, b or full protein
- $[^3\text{H}]$ thymidine incorporation assay


T cells respond to immunization with isoAsp self-peptide and fail to respond Asp self-peptide

The effect of post-translational modification on T-cell immune recognition: Isomerisation of Asp to β -Asp


peptide 94-104 from murine cytochrome c



NMR analysis of immunogenic/non-immunogenic self-peptide










$^{94}\text{ERADLIAYLKKTNE}^{104}$







$^{94}\text{ERAIsoDLIAYLKKTNE}^{104}$

The effect of post-translational
modification on antibody recognition:
glycosylation


Glycosylation




hydroxylation







Identification of antibody epitope of mucin-2







Healthy tissue




Tumor tissue



The effect of carbohydrate moiety on MoAb binding





The effect of carbohydrate moiety on MoAb binding



MAb 996:


Anti-BSA-[K¹²VTPPTPTGTQTPT²⁵]

The effect of carbohydrate moiety on MoAb binding




IC₅₀ (μM) = > 1000

The effect of carbohydrate moiety on MoAb binding



$^{16}\text{Pro-Thr-Pro-Thr-Gly-Thr-Gln}^{22}$


$\text{IC}_{50} (\mu\text{M}) = 25$



$^{16}\text{Pro-Thr-Pro-Thr-Gly-Thr-Gln}^{22}$


$\text{IC}_{50} (\mu\text{M}) = 39$

The effect of carbohydrate moiety on MoAb binding




$^{16}\text{Pro-Thr-Pro-Thr-Gly-Thr-Gln}^{22}$

$\text{IC}_{50} (\mu\text{M}) = 25$




$\text{IC}_{50} (\mu\text{M}) = 6.4$

$^{16}\text{Pro-Thr-Pro-Thr-Gly-Thr-Gln}^{22}$




The effect of carbohydrate moiety on MoAb binding




$^{16}\text{Pro-Thr-Pro-Thr-Gly-Thr-Gln}^{22}$

$IC_{50} (\mu\text{M}) = 25$




$IC_{50} (\mu\text{M}) = 6.4$

$^{16}\text{Pro-Thr-Pro-Thr-Gly-Thr-Gln}^{22}$



$^{16}\text{Pro-Thr-Pro-Thr-Gly-Thr-Gln}^{22}$


$IC_{50} (\mu\text{M}) = 39$



$^{16}\text{Pro-Thr-Pro-Thr-Gly-Thr-Gln}^{22}$


$IC_{50} (\mu\text{M}) = > 1000$

The effect of carbohydrate moiety on MoAb binding




$^{16}\text{Pro-Thr-Pro-Thr-Gly-Thr-Gln}^{22}$

$IC_{50} (\mu\text{M}) = 25$




$IC_{50} (\mu\text{M}) = 6.4$

$^{16}\text{Pro-Thr-Pro-Thr-Gly-Thr-Gln}^{22}$




$^{16}\text{Pro-Thr-Pro-Thr-Gly-Thr-Gln}^{22}$

$IC_{50} (\mu\text{M}) = 39$



$IC_{50} (\mu\text{M}) = > 1000$



$IC_{50} (\mu\text{M}) = > 1000$

ECD (Electronic Circular Dichroism)

Instrument: Jasco-810

Solvent:

- water
- TFE




<http://www.andrew.cmu.edu/user/jamess3/JWSfac.html>

Concentration: 0,5 mg/ml


Wavelength: $\lambda=180-300$ nm

0,02 cm quartz cuvette

The circular dichroism spectra of peptides in TFE










The circular dichroism spectra of peptides in water







The effect of post-translational
modification on antibody recognition:
citrullination


Citrullination




hydroxylation



Rheumatoid arthritis




Pierre Auguste Renoir (1841 - 1919)



**Dorothy C. Hodgkin (1910-1994)
Nobel dij (1964)**



Raoul Dufy (1877-1953)





Post-translational modification: Immune recognition related diseases

Modification	Autoantigen	Disease
Acetylation	Myelin basic protein	Multiple sclerosis
Citrullination	Collagen type II Myelin basic protein	RA Multiple sclerosis
Deamidation	Insulin	Type I diabetes
Glycosylation	Insulin proceptor Collagen type II Thyrotropin receptor Myelin oligodendrocyte glycoprotein	Diabetes RA Graves disease MS
Isoaspartylation	snRNP	Systemic lupus erythematosus
Lipoylation	PDC-E2	Primary biliary cirrhosis
Phosphorylation	Myelin basic protein	Multiple sclerosis
Methylation	Sm, D1,D3	Systemic lupus erythematosus
Transglutamination	Histone H2	Systemic lupus erythematosus
Tyrosine nitration	Mitochondrial proteins	Experimental autoimmune uveitis

Rheumatoid Arthritis

- chronic, systemic inflammatory disorder
- systemic autoimmune disease
- attacks synovial joints
- hyperplasia of synovial cells,
- excess synovial fluid,






http://en.wikipedia.org/wiki/File:Rheumatoid_arthritis_world_map_-_DALY_-_WHO2004.svg



Epidemiology


- 1% of the world's population
- women : man = 3:1
- most frequent ages 40 - 50

The effect of post-translational modification on immune recognition: change in 3D structure of proteins



Yamada, R. et al. *Bioscience* 10: 54-64 (2005)
 Yamada, R. *Autoimmunity Reviews* 4: 201-206 (2005)

Immunity to citrullinated proteins in rheumatoid arthritis



The effect of post-translational modification on immune recognition: proteins involved


Disease	Modification	Antigen modified
RA	Hydroxylation	Type II collagen
	Glycosylation	
	Oxidation	<div style="border: 1px solid red; padding: 2px;"> Fillagrin Fibrin Vimentin </div>
	Citrullination	
	Glycosylation	IgG
SLE	Phosphorylation	Multiple snRNP D, H2B
	Deamidation	
	Mannose modification	Multiple SM D1, D3
	Methylation	
	Oxidation	Cardiolipin, ox LDL, C1q, calreticulin

Eggleton, P. et al. *Rheumatology* 47: 567-571 (2008)




filaggrin¹

vimentin²



Based on [PyMOL](#) rendering of PDB [1gk4](#)



fibrin³

- 1 Sebbag, M. et al. *Clin. Invest.* 95: 2672-2679 (1995)
- 2 Vossenaar, E.R. et al. *Arthritis Res. Ther.* 6(2): 86-89 (2004)
- 3 Masson-Bessiere, C. et al. *J. Immun.* 166: 4177-4184 (2001)


Crystal Structure of Fibrin from *Petromyzon marinus*, [1n73](#)
<http://www.proteopedia.com/wiki/index.php/Image:1n73b.png>

The effect of post-translational modification on immune recognition: **epitope peptide**


Vimentin	⁶⁵ SAV RAR SSVPGV RK ⁷⁷
Fibrin α	³⁴ GPRVV R HQSACKDS ⁴⁸
Fibrin β	⁶⁰ RPAPPPISSGGY RAR ⁷⁴
Filaggrin (5-mer)	³¹¹ TR GR S ³¹⁵
Filaggrin (19-mer)	³¹¹ SHQEST RGR SRGRSGRSGS ³²⁶



filaggrin



vimentin



fibrin ¹

¹ Iobagiu C., Magyar, A. et al. *J. Autoimmunity* 37: 263-272 (2011)

Aims

1. Identification of minimal and optimal antibody epitope of partially deimidated filaggrin by synthetic peptides based on 306-324 sequence using multi-pin approach and serum samples from diseased individuals.
2. Introduction of biotin label for soluble epitope peptide
3. Analyze
 - the effect epitope size and orientation on antibody recognition,
 - the effect the presence and position of biotin on solution conformation,
 - RA specificity in serum samples as compared with that of SLE and healthy individuals using the optimized peptide epitope by direct ELISA.

Aims

1. Identification of minimal and optimal antibody epitope of partially deimidated filaggrin by synthetic peptides based on 306-324 sequence using multi-pin approach and serum samples from diseased individuals.
2. Introduction of biotin label for soluble epitope peptide
3. Analyze
 - the effect epitope size and orientation on antibody recognition,
 - the effect the presence and position of biotin on solution conformation,
 - RA specificity in serum samples as compared with that of SLE and healthy individuals using the optimized peptide epitope by direct ELISA.

Filaggrin (filament aggregating protein)

(FILA_HUMAN), <http://swissmodel.expasy.org/>

profilaggrin: 4061 AA, 435170 Da





10-12 filaggrin unit, 324 AA

SHQESTRGRSRGRSGRSGS *

MSTLLENIFA IINLFKQYSK KDKNTDLSK KELKELLEKE FRQILKNPDD PDMVDVFMH LDIDHNKID FTEFLLMVFK LAQAYYESTR KENLPISGHK HRKHSHHDKH EDNKQEENKE NRKRPSSLER RNNRKGNGKR SKSPRETGGK RHESSSEKKE RKGYSPTRE EYKGNHNS SKKEKNKTEN TRLGDNKRRL SERLEEKEDN EEGVDYENT GRMTQKWIQS GHIAITYYTIQ DEAYDTTDSL LEENKIYERS RSSDQKSSSQ VNRSRHENTS QVPLQESRTR KRRGSRVSDQ RDEGHSSEDS ERHSGSASRN HHGSAWEQSR DGSRRHPRSHD EDRASHGHS A DSSRQSGTRH AETSSRGQTA SSHEQARSSP GERHSGSHQO SADSSRHSAT GRGQASSAVS DRGHRGSSGS QASDSEGHSE NSDTQSVSGH GKAGLRQQSH QESTRGRSGE RSGRSGSSLY QVSTHEQPDS AHGRTGTSTG GRQGSHEQA RDSSRHSAQ EGQDTIRGHP GSSRGGRQGS HHEQSVNRSR HSGSHHSHTT SQGRSDASHG QSGRSASRQ TRNEEQSGDG TRHSGSRHHE ASSQADSSRH SQVGQGSQSSG PRTSRNQSS VSQSDSQGH SEDSERWSGS ASRNHHGSAQ EQSRDGRHP RSHHEDRAGH GHSADSSRKS GTRHTQNSSS GQAASSHEQA RSSAGERHGS RHQLQSADSS RHSGTGHGQA SSAVRDSGHR GSSGSQATDS EGHSESDTQ SVSGHGQAGH HQQSHQESAR DRSGERSRRS GSFLYQVSTH KQESSHGWT GPSTGVRQGS HHEQARDNSR HSASQDQDQD IRGHPGSSRR GRQGSHEQS VDRSGHSGSH HSHTTSQGRS DASRGQSGSR SASRTRNEE QSRDGRHSG SRHHEASSHA DISRHSQAGQ GQSEGSRTSR RQGSVSDS DSEGHSEDE RWSGSASRNH RGSAQEQSRH GSRHPRSHHE DRAGHGSAD SSRQSGTPHA ETSSGGQAAS SHEQARSSP ERHGSRHQQS ADSSRHSGIP RRQASSAVRD SGHWGSSGSQ ASDSEGHSEE SDTQSVSGHG QDQDPHQSSHQ ESARDWSGGR SGRSGSFTYQ VSTHEQSESA HGRTRTSTGR RQGSHHEQAR DSSRHSAQSE QDRTIRAHPG SRRRGGQGSHEQSVDRSGH GSHHSHTT QGRSDASHGQ SGRSASRQT RKDKQSGDGS RHSGSRHHEA ASWADSSRHS QVQEQSSGS RTSRHQGSVV SQDSDSERHS DDSERLGSQA SRNHHGSSRE QSRDGRHPG FHQEDRASHG HSADSSRQSG THHTESSHG QAVSSHEQAR SSPGERHGSR HQQSADSSRH SGIGHRQASS AVRDSGHRGS SGSQVTNSEG HSESDTQSV SAHQAGPHQ QSHKESARGQ SGESSGRSRS FLYQVSSHEQ SESTHGQTAP STGGRQGSRH EQARNSSRHS ASQDQDQD GHPGSSRGGR QGSYHEQSVDR RSGHSGYHHS HTTPQGRSDA SHGQSGPRSA SRQTRNEEQS GDGSRHSGSR HHEPSTRAGS SRHSQVQGE SAGSKTSRRQ GSSVQDRDS EGHSEDSERR SESASRNHYG SAREQSRHGS RNPRSHQEDR ASHGSAESS RQSGTRHAET SGGQAASSQ EQARSSPGER HGSRHQQSAD SSTDSGTGRR QDSSVWDSG NRGSSGSQAS DSEGHSEED TQSVSAHGQA GPHQSHQES TRGQSGERSG RSGSFLYQVS THEQSESAHG RTGPSTGGRQ RSRHEQARDS SRHSASQEGQ DTIRGHPGSS RGGRQGSHEQ QSVDSGSHG SHHSHTTSQE RSDVSRGQSG SRSVSRQTRN EKQSGDGRH SGRHHEASS RADSSRHSQV GQGSQSGPRT SRNQSSVSQ DSDSQGHSE SERWSGSASR NHLGSAWEQS RDGSRHPGSH HEDRAGHGH ASSSRQSGTR HTESSSRQA ASSHEQARSS AGERHSGHHQ LQADSSRHS GIGHGQASSA VRDSGHRGYS GSQASDSEGH SEDSDTQSVS AQQKAGPHQ SHKESARGQS GESSRSGSF LYQVSTHEQS ESTHGQSAPS TGGRQGSYD QAQDSSRHS SQEGQDTIRG HPGPSRGGRQ GSHQEQSVDR SGHSGSHHS TTSQGRSDAS RGQSGRSAS RKTIDYKEQG DGRHSGSHH HEASSWADSS RHLVQGQGS SGPRTSRPRG SSVQSDSDSE GHSEDSERRS GSASRNHHS AQEQSRDGRS HPRSHHEDRA GHGSAESSR QSGTHHAENS SGGQAASSHE QARSSAGERH GSHHQQSADS SRHSGIGHGQ ASSAVRDSGH RGSQSGASD SEGHSESDT QSVSAHGQAG PHQSHQEST RGRSAGRSGR SSGSFLYQVST HEQSESAHGR TGTSTGGRQ SHHKQARDSS RHSTSQEQD TIHGHPGSSS GGRQGSHEQ LVDRSGHSGS HHSHTTSQGR SDASHGHSGS RSASRQTRND EQSGDGRHS GSRHHEASSR ADSSGHSQV QGQSEGPRTS RNWGSFSQD SDSQGHSEDS ERWSGSASRN HHGSAQEQLR DGRHPRSHQ EDRAGHGHSA DSSRQSGTRH TQTSSGGQA SSHEQARSSA GERHSGHHQ SADSSRHSGI GHGQASSAVR DSGHRGYSGS QASDNFEGHSE DSDTQSVSAH GQAGSHQQS QESARGRSGE TSGHSGSFLY QVSTHEQSES SHGWTGPSTR GRQGSRHEQA QDSSRHSAQ DGQDTIRGHP GSSRGGRQGS HHEHSDSSG HSGSHHSHTT SQGRSDASRG QSGRSASRT TRNEEQSGDG SRHSGSRHHE ASTHADISRH SQAVQGSQEG SRRSRQGS VSQSDSEGH SEDSERWSGS ASRNHHGSAQ EQLRDSGRHP RSHQEDRAGH GHSADSSRKS GTRHTQTSSG GQAASSHEQA RSSAGERHGS HHQQSADSSR HSGIGHGQAS SAVRDSGHRG YSGSQASDNE GHSESDTQS VSAHQAGSH QQSHQESARG RSGETSGHSG SFLYQVSTHE QESSHGWTG PSTRGRQGSR HEQAQDSSRH SASQYQDQTI RGHGSSRGG RQGYHHEHSV DSSGHSGSHH SHTTSQGRSD ASRGQSGRS ASRTRNEEQ SGDSSRHVS RHHEASTHAD ISRHSQAVQG QSEGSRRSR RQGSVSDSD SEGHSEDSER WSGSASRNHR GSVQEQSRHG SRHPRSHHD RAGHGHSA DR SRQSGTRHAE TSSGGQAASS HEQARSSPGE RHGSRHQQA DSSRHSGIPR GQASSAVRDS RHWSGSSQA SDSEGHSES DTQSVSGHGQ AGPHQSHQ SARDRSGRS GRSGSFLYQV STHEQSESAH GRTRTSTGR QGSHHEQARD SSRHSASQEG QDTRGHPGS SRRRQGSHEQ SVDRSGHS GSHHSHTTSQ GRSDASRGQS GRSASRQTR NDEQSGDGR HSWSHHEAS TQADSSRHSG QDQGSAGPR TSNRQGSVS QDSDSGHSE DSERWGSAS RNHRGSAQEQ SRDGRHPTS HHEDRAGHGH SAESSRQSGT HHAENSSGGQ AASSHEQARS SAGERHSGSH QQSADSSRH GIGHGQASSA VRDSGHRGSS GSQASDSEGH SEDSDTQSVS AHGQAGPHQ SHQESTRGRS RGRSGRSGS LYQVSTHEQS ESAHGRAGPS TGGRQGSRHE QARDSSRHS SQEGQDTIRG HPGSRRGGRQ GSYHEQSVDR SGHSGSHHS TTSQGRSDAS HGQSGRSAS RETRNEEQSG DGRHSGSRH HEASTQADSS RQSQSGQES AGSRRSRQGS SSVQSDSDSE AYPEDSERRS ESASRNHHS SREQSRDGR HPGSSHRDTA SHVQSSPVQS DSSTAKEHGH FSSLQDSAY HSGIQSRGSP HSSSYHYQS EGTERQKQGS HGVS

* Schellekens, G.A. et al. *J. Clin. Invest.* 101, 273-281 (1998)




Search for minimal/optimal epitope: Multi-pin approach



Deprotection

- 1) 20 % piperidine/DMF
- 2) TFA/tioanisole/anisole/water/EDT =
= 82,5 : 5 : 5 : 5 : 2,5 (v/v/v/v/v)

Identification of linear antibody epitopes



In vitro analysis of antibody recognition


- Peptides
 - C-terminally pin-bound oligopeptides
 - free peptides labelled with biotin at the N-terminal
 - free peptides labelled with biotin at the C-terminal
- Serum samples:
 - from healthy CCP positive individuals,
 - from diseased CCP positive RA individuals,
 - from healthy CCP negative individuals,
- Synovial fluid samples:
- direct ELISA

Search for minimal/optimal epitope

19-mer analogues	N-terminal truncation of peptide Cit ³¹² (306-324)	C-terminal truncation of peptide 311-324
³⁰⁶ SHQESTRGRSRGRSGRSGS ³²⁴	³⁰⁶ SHQESTXGRSRGRSGRSGS ³²⁴	³¹¹ TXGRSRGRSGRSGS ³²⁴
³⁰⁶ SHQESTXGXSGXSGXSGS ³²⁴	³⁰⁷ HQESTXGRSRGRSGRSGS ³²⁴	³¹¹ TXGRSRGRSGRSG ³²³
³⁰⁶ SHQESTXGXSGXSGRSGS ³²⁴	³⁰⁸ QESTXGRSRGRSGRSGS ³²⁴	³¹¹ TXGRSRGRSGRS ³²²
³⁰⁶ SHQESTXGXSGRSGRSGS ³²⁴	³⁰⁹ ESTXGRSRGRSGRSGS ³²⁴	³¹¹ TXGRSRGRSGR ³²¹
³⁰⁶ SHQESTXGXSRGRSGRSGS ³²⁴	³¹⁰ STXGRSRGRSGRSGS ³²⁴	³¹¹ TXGRSRGRSG ³²⁰
³⁰⁶ SHQESTXGRSRGRSGRSGS ³²⁴	³¹¹ TXGRSRGRSGRSGS ³²⁴	³¹¹ TXGRSRGRS ³¹⁹
³⁰⁶ SHQESTRGXSGXSGXSGS ³²⁴	³¹² XGRSRGRSGRSGS ³²⁴	³¹¹ TXGRSRGR ³¹⁸
³⁰⁶ SHQESTRGRSXXSGXSGS ³²⁴		³¹¹ TXGRSRG ³¹⁷
³⁰⁶ SHQESTRGRSRGXSGXSGS ³²⁴		³¹¹ TXGRSR ³¹⁶
³⁰⁶ SHQESTRGRSRGRSGXSGS ³²⁴	Control peptides	³¹¹ TXGRS ³¹⁵
³⁰⁶ SHQESTXGXSRGRSGRSGS ³²⁴	PLAQGGGGGG	³¹¹ TXGR ³¹⁴
³⁰⁶ SHQESTXGRSXXGRSGRSGS ³²⁴	GLAQGGGGGG	³¹¹ TRGR ³¹⁴
³⁰⁶ SHQESTXGRSRGRSGXSGS ³²⁴		


(X=citrullin)

Search for minimal/optimal epitope



Magyar, A. et al. in *Peptides 2000, Proc. 26th European Peptide Symposium* (Ed.: Martinez, J., Fehrentz, J-A.) EDK, Paris, France, 679-680 (2001)

Analogue 19-mer peptides: Critical Cit residue(s)




19-mer analogues
306 SHQESTRGRSRGRSGRSGS³²⁴ *
306 SHQESTXGXSGXSGXSGS ³²⁴
306 SHQESTXGXSGXSGRSGS ³²⁴
306 SHQESTXGXSGRSGRSGS³²⁴
306 SHQESTXGXSRGRSGRSGS ³²⁴
306 SHQESTXGRSRGRSGRSGS ³²⁴
306 SHQESTRGRSRGXSGXSGS ³²⁴
306 SHQESTRGRSRGXSGXSGS ³²⁴
306 SHQESTRGRSRGRSGXSGS ³²⁴
306 SHQESTXGXSRGRSGRSGS ³²⁴
306 SHQESTXGRSRGXGRSGRSGS ³²⁴
306 SHQESTXGRSRGRSGXSGS ³²⁴

(X = citrullin)

*based on Hu-profilaggrin cDNS aa 306-324


N-terminal truncation of peptide Cit³¹² (306-324)



N-terminal truncation of peptide Cit ³¹² (306-324)	
306	SHQESTXGRSRGRSGRSGS ³²⁴
307	HQESTXGRSRGRSGRSGS ³²⁴
308	QESTXGRSRGRSGRSGS ³²⁴
309	ESTXGRSRGRSGRSGS ³²⁴
310	STXGRSRGRSGRSGS ³²⁴
311	TXGRSRGRSGRSGS ³²⁴
312	XGRSRGRSGRSGS ³²⁴

(X = citrullin)

C-terminal truncation of peptide Cit³¹² (311-324)





C-terminal truncation of peptide 311-324	
311	TXGRSRGRSGRSGS ³²⁴
311	TXGRSRGRSGRSG ³²³
311	TXGRSRGRSGRS ³²²
311	TXGRSRGRSGR ³²¹
311	TXGRSRGRSG ³²⁰
311	TXGRSRGRS ³¹⁹
311	TXGRSRGR ³¹⁸
311	TXGRSRG ³¹⁷
311	TXGRSR ³¹⁶
311	TXGRS ³¹⁵
311	TXGR ³¹⁴
311	TRGR ³¹⁴

(X = citrullin)


Aims

1. Identification of minimal and optimal antibody epitope of partially deimidated filaggrin by synthetic peptides based on 306-324 sequence using multi-pin approach and serum samples from diseased individuals.
2. Introduction of biotin label for soluble epitope peptide
3. Analyze
 - the effect epitope size and orientation on antibody recognition,
 - the effect the presence and position of biotin on solution conformation,
 - RA specificity in serum samples as compared with that of SLE and healthy individuals using the optimized peptide epitope by direct ELISA.


Analysis of antibody binding to biot- β 60-74Cit/Arg peptide by direct ELISA





The effect of epitope orientation: the position of biotin




The effect of epitope accessibility: linkers



Biotinyl-4,7,10-trioxa-1,13-tridecandiamino-succinic acid



Synthesis of biotinyl-4,7,10-trioxa-1,13-tridecandiamino-succinic acid



Bartos, Á. et al.
Biopolymers 92: 110-115 (2009)
Bartos, Á. et al.
Tetrahedron Letters 50: 2661-2663 (2009)

Aims

1. Identification of minimal and optimal antibody epitope of partially deimidated filaggrin by synthetic peptides based on 306-324 sequence using multi-pin approach and serum samples from diseased individuals.
2. Introduction of biotin label for soluble epitope peptide
3. Analyze
 - the effect epitope size and orientation on antibody recognition,
 - the effect the presence and position of biotin on solution conformation,
 - RA specificity in serum samples as compared with that of SLE and healthy individuals using the optimized peptide epitope by direct ELISA.

Analysis of serum samples

Samples:

- 263 RA patients with established disease,
- 46 CCP negative, non-RA patients with other autoimmune diseases
- 18 patients with systemic lupus erythematosus
- 152 age-matched healthy controls

The diagnosis of the disease was established on the basis of the revised ACR/EULAR classification criteria. ¹


The baseline data of RA patients:

32 men/176 women; age: 58,4 +/- 14,3 years;
rheumatoid factor (RF) +/-: 127/30; CCP2 +/-: 157/27; MCV +/-: 164/25;
disease duration: 9.8 +/- 9,4 years.


Statistical analysis: ANOVA, compared with Pearson's correlation analysis

1. Aletaha, D., Neogi, T., Silman, A.J. et al. 2010 rheumatoid arthritis classification criteria: an American College of Rheumatology/European League Against Rheumatism collaborative initiative. *Ann Rheum Dis* 69:1580-8 (2010).


Synthesis of 5-mer epitope peptide with *N*-terminal biotin




biotinyl-TXGRS-NH₂



biotinyl-6-aminohexanoyl-TXGRS-NH₂




biotinyl-TRGRS-NH₂



biotinyl-6-aminohexanoyl-TRGRS-NH₂


Compound	M_{av} (calc)	M_{av} (meas)	R_t (min)
biotinyl-TRGRS-NH ₂	802,9	802,8	14,23
biotinyl-TXGRS-NH ₂	803,9	803,7	14,12
biotinyl-6-aminohexanoyl-TRGRS-NH ₂	914,1	913,9	17,07
biotinyl-6-aminohexanoyl-TXGRS-NH ₂	915,1	914,9	16,50

Antibody recognition of 5-mer epitope peptides with *N*-terminal biotin




$$OD \text{ ratio} = OD_{\text{Cit peptide}} / OD_{\text{unmodified peptide}}$$

Synthesis of 5-mer epitope peptides with C-terminal biotin




Ac-TRGRSK(biotinyl-hexanoyl)-NH₂




Ac-TRGRS-Ttds-K(biotinyl-hexanoyl)-NH₂

Ac-TRGRSK(biotinyl-Ttds)-NH₂



Ac-TXGRSK(biotinyl-hexanoyl)-NH₂



Ac-TXGRS-Ttds-K(biotinyl-hexanoyl)-NH₂

Ac-TXGRSK(biotinyl-Ttds)-NH₂


Characteristics of 5-mer epitope peptide with C-terminal biotin

Compound	M _{av} (calc)	M _{av} (meas)	R _t (min)
Ac-TRGRSK(biotinyl-hexanoyl)-NH ₂	1084,3	1084,6	17,08
Ac-TXGRSK(biotinyl-hexanoyl)-NH ₂	1085,3	1084,6	17,05
Ac-TRGRSK(biotinyl-Ttds)-NH ₂	1273,1	1273,2	17,97
Ac-TXGRSK(biotinyl-Ttds)-NH ₂	1274,1	1273,9	17,62
Ac-TRGRS-Ttds-K(biotinyl-hexanoyl)-NH ₂	1386,3	1386,4	18,16
Ac-TXGRS-Ttds-K(biotinyl-hexanoyl)-NH ₂	1387,3	1387,4	18,42


HPLC: KNAUER, Synergi MAX-RP, C12, 250 x 4mm, 5µm silica, 100 Å column, 5%B - 95 % B, 50 min,
eluent A: 0,1% TFA/water (V/V); eluent B: 0,1% TFA/acetonitrile-water (80:20 V/V)

MS: Esquire 3000+


Antibody recognition of 5-mer epitope peptides with C-terminal biotin



Synthesis of 19-mer epitope peptide (³⁰⁶SHQESTRGRSRGRSGRSGS³²⁴) with *N*-terminal biotin





biotinyl-6-aminohexanoyl-SHQESTRGRSRGRSGRSGS-NH₂



biotinyl-6-aminohexanoyl-SHQESTXGXSGRSGRSGS-NH₂

Compounds	M _{av} (calc)	M _{av} (meas)	R _t (min)
biotinyl-6-aminohexanoyl-SHQESTRGRSRGRSGRSGS-NH ₂	2383,6	2383,8	13,27
biotinyl-6-aminohexanoyl-SHQESTXGXSGRSGRSGS-NH ₂	2386,6	2386,7	12,95


Antibody recognition of 19-mer epitope peptide with *N*-terminal biotin by RA, SLE and healthy samples




The presence of biotin at *N*-terminal biotin
does not prevent recognition.

RA vs SLE	P < 0.001
RA vs Healthy	P > 0.05
SLE vs Healthy	P > 0.05


Synthesis of 19-mer epitope peptide (³⁰⁶SHQESTRGRSRGRSGRSGS³²⁴) with C-terminal biotin



Ac-SHQESTRGRSRGRSGRSGS-Ttds-K(biotinyl-6-aminohexanoyl)-NH₂



Ac-SHQESTRGRSRGRSGRSGSK(biotinyl-6-aminohexanoyl)-NH₂



Ac-SHQESTRGRSRGRSGRSGSK(biotinyl-Ttds)-NH₂



Characteristics of 19-mer epitope peptide (³⁰⁶SHQESTRGRSRGRSGRSGS³²⁴) with C-terminal biotin

Compound	M _{av} (calc)	M _{av} (meas)	R _t (min)
Ac-SHQESTRGRSRGRSGRSGSK(biotinyl-aminohexanoyl)-NH ₂	2553,8	2553,8	14,23
Ac-SHQESTXGXSGRSGRSGSK(biotinyl-aminohexanoyl)-NH ₂	2556,8	2556,9	13,73
Ac-SHQESTRGRSRGRSGRSGSK(biotinyl-Ttds)-NH ₂	2743,3	2743,4	15,25
Ac-SHQESTXGXSGRSGRSGSK(biotinyl-Ttds)-NH ₂	2746,3	2746,5	14,90
Ac-SHQESTRGRSRGRSGRSGS-Ttds-K(biotinyl-aminohexanoyl)-NH ₂	2856,4	2856,5	17,65
Ac-SHQESTXGXSGRSGRSGS-Ttds-K(biotinyl-aminohexanoyl)-NH ₂	2859,4	2859,5	17,35

HPLC: KNAUER, Synergi MAX-RP, C12, 250 x 4mm, 5µm silica, 100 Å column, 5%B - 95 % B, 50 min, eluent A: 0,1% TFA/water (V/V); eluent B: 0,1% TFA/acetonitrile-water (80:20 V/V)

MS: Esquire 3000+

Antibody recognition of 19-mer epitope peptide with C-terminal biotin by RA, SLE and healthy samples




Ac-SHQESTXGXSRSGSGSK(biotinyl-6-aminohexanoyl)-NH₂

The peptide with C-terminal biotin
is recognized.

RA vs SLE	P < 0.001
RA vs Healthy	P > 0.05
SLE vs Healthy	P > 0.05


Antibody recognition of 19-mer epitope peptide with *N*- or *C*- terminal biotin

biotinyl-6-aminohexanoyl-
-SHQESTXGXSGRSGRSGS-NH₂



RA vs SLE		P < 0.001
RA vs Healthy		P > 0.05
SLE vs Healthy		P > 0.05

Ac-SHQESTXGXSGRSGRSGSK
(biotinyl-aminohexanoyl)-NH₂




RA vs SLE		P < 0.001
RA vs Healthy		P < 0.01
SLE vs Healthy		P < 0.05

Both *C*- and *N*-terminal biotinylated 19-mer epitope peptides are recognized by RA sera samples

Comparison of antibody recognition of the 5-mer and the 19-mer epitope peptide

OD ratio (citrullinated/non-citrullinated)


Filaggrin19



RA vs CCP neg.	P < 0.001
RA vs Healthy	P < 0.001
CCP neg. vs Healthy	P > 0.05

OD ratio (citrullinated/non-citrullinated)

Filaggrin 5







RA vs CCP neg.	P < 0.001
RA vs Healthy	P < 0.001
CCP neg. vs Healthy	P > 0.05

Short summary

N-terminal biotinylation

C-terminal biotinylation



ECD (Electronic Circular Dichroism)

Instrument: Jasco-810

Solvent:

- water
- TFE




<http://www.andrew.cmu.edu/user/jamess3/JWSfac.html>

Concentration: 0,5 mg/ml


Wavelength: $\lambda=180-300$ nm

0,02 cm quartz cuvette


The effect of biotin position on ECD spectra of 19-mer in water



The effect of biotin position on ECD spectra of 19-mer in TFE



The position of the epitope core within the epitope region influence the antibody recognition



Conclusions


1. An epitope region (19-mer) and an epitope core (5-mer) were identified.
2. Introduction of biotin to the *N*-terminal of the 5-mer resulted in **no binding**. The presence of biotin at the *C*-terminal of the 5-mer had no effect on binding.
3. However, the **presence of biotin at *N*- or *C*-terminal of the 19-mer has no effect on epitope recognition** by serum antibodies.
4. **The position of biotin markedly influences the solution conformation of the epitope peptide (5-mer)**. In contrast, no influence of the biotin position could be detected in case of 19-mer epitope region peptide.
5. The 5-mer as well as the 19-mer citrullinated peptides have shown a **significantly higher reactivity with CCP^+ RA sera as compared to healthy controls, CCP^- serum samples**.

Acknowledgements

F. Babos¹, E. Szarka², A. Magyar¹, G. Sármay²

¹ Research Group of Peptide Chem., Hungarian Academy of Sciences, Eötvös L. University,

² Department of Immunology, Eötvös L. University, Budapest



Pázmány Péter program



CellKom

Regionális Egyetemi Tudásközpont



Pázmány

A projekt a Nemzeti Kutatási és Technológiai Hivatal támogatásával valósult meg.



Support

Hungarian-French Intergovernmental Program (F-9/2010)

Hungarian Academy of Sciences

Hungarian National Research Fund (OTKA T045634)

Ministry of Health (ETT)